





The light-curing bracket adhesive.

For the adhesion of metal, ceramic and plastic brackets and buccal tubes.

CONTEC lc offers all the advantages of a light-curing adhesive. This gives you more time to place the brackets before the adhesive cures.

The high viscosity ensures that the adhesive stays on the bracket and does not drift. At the same time, the bond strength of CONTEC lc is excellent. It is therefore the solution for all bonding tasks.

The ergonomically-formed syringe makes for easy handling and good control over the amount of adhesive used.

The special advantage of CONTEC lc is that it is free of BisGMA*, TEGDMA* and HEMA*. These materials have been proven to have allergenic potential.



CONTEC lc set (REF 163-110-00) consisting of:

O 3 x 3 g CONTEC lc adhesive

O 1 x 5 ml CONTEC lc/lcr primer

O 2 x 2.5 ml CONTEC etching gel

O 6 x CONTEC disposable tube for etching gel

O 50 x CONTEC disposable brush

O 1 x CONTEC combination instrument

O 1 x CONTEC mixing pad

For approx. 800 brackets (40 treatment cases).



* CONTEC lc contains no BisGMA (bisphenol A glycerolate dimethacrylate), TEGDMA (triethylene glycol dimethacrylate) or HEMA (hydroxyethyl methacrylate).

CONTEC Ic



The light-curing retainer adhesive.

For the adhesion of fixed retainers.



CONTEC lcr has been developed to fulfill the special requirements for bonding fixed orthodontic retainers. With CONTEC lcr, retainers can be secured safely and with the maximum patient comfort possible. Yet the adhesive is durable and lasting and has excellent bonding characteristics.

The slightly bent tubes and the delicate syringes make handling in the lingual area of the anterior teeth easy.

CONTEC lcr is free of TEGDMA* and HEMA*. CONTEC lcr is therefore the first choice for patients who have been diagnosed with an allergy against one of these substances.

Single pack	Contents	REF
CONTEC lcr adhesive	1 x 3.4 g syringe	163-131-00
CONTEC lc/lcr primer	1 x 5 ml bottle	163-113-10
CONTEC etching gel	2 x 2.5 ml syringe	163-115-00

Developed in cooperation with:

SAREMCO

* CONTEC lcr is free of TEGDMA (triethylene glycol dimethacrylate) and HEMA (hydroxyethyl methacrylate).

CONTE Icr



The ready-to-use, chemically-curing adhesive.

For adhering metal, ceramic and plastic brackets and buccal tubes.



CONTEC go is a chemically curing, ready-to-use bracket adhesive consisting of an adhesive paste and a primer. The curing process starts by means of pressure. This allows the brackets to be positioned on the tooth before the adhesive cures.

There is sufficient strength in the bond after only 5 minutes for the archwire to be ligated.


CONTEC go has a high viscosity and a consistency that guarantees optimum stability on the bracket. CONTEC go also maintains its color throughout the period of treatment.

CONTEC go is free of TEGDMA* and HEMA*.

CONTEC go is therefore the first choice for patients who have been diagnosed with an allergy against one of these substances.

Single pack	Contents	REF
CONTEC go adhesive	2 x 2.5 g syringe	163-121-00
CONTEC go primer	1 x 5 ml bottle	163-114-00
CONTEC etching gel	2 x 2.5 ml syringe	163-115-00





Dentaurum

Germany I Benelux I España I France I Italia I Switzerland I Australia I Canada I USA and in more than 130 countries worldwide.



DENTAURUM QUALITY WORLDWIDE UNIQUE





Date: 2022-10 Subject to modifications